

Scaling Relations and Exponents in the Growth of Rough Interfaces Through Random Media

J.M. López,⁽¹⁾ M.A. Rodríguez,⁽¹⁾ A. Hernández-Machado,⁽²⁾ and A. Díaz-Guilera⁽³⁾

⁽¹⁾*Departamento de Física Moderna, Universidad de Cantabria, and Instituto de Estudios Avanzados en Física Moderna y Biología Molecular, CSIC-UC, Avenida Los Castros, E-39005 Santander, Spain.*

⁽²⁾*Departament d'Estructura i Constituents de la Matèria, Universitat de Barcelona, Diagonal 647, E-08028 Barcelona, Spain.*

⁽³⁾*Departament de Física Fonamental, Universitat de Barcelona, Diagonal 647, E-08028 Barcelona, Spain.*

The growth of a rough interface through a random media is modelled by a continuous stochastic equation with a quenched noise. By use of the Novikov theorem we can transform the dependence of the noise on the interface height into an effective temporal correlation for different regimes of the evolution of the interface. The exponents characterizing the roughness of the interface can thus be computed by simple scaling arguments showing a good agreement with recent experiments and numerical simulations.

05.40.75, 05.60c, 05.70L

The growth of rough interfaces through a random medium is an interesting pattern-formation phenomena that has received a lot of attention in recent years [1–12]. In particular, experimental studies of the displacement of a nonwetting fluid by a more viscous wetting one through a porous media have shown that the interface separating both fluids is rough and the random pinning of the displacing fluid on the wider porous of the medium strongly influences the growth [2,5]. The random media is frozen in time and a natural way to model its effects is by introducing a quenched disorder in a continuous equation, as has been done by Kessler, Levine and Tu (KLT) [3], Parisi [7], and others [6,10]. This model is different from the one that describes the growth of solids from a vapor or from a stochastic ballistic deposition, in which the stochastic field changes both in space and time. These last phenomena are usually described by equations of the type introduced by Edwards and Wilkinson (EW) [13] and by Kardar, Parisi and Zhang (KPZ) [14]. The effects of a quenched noise in a stochastic model are much less known and are the subject of this Letter.

We study the model introduced by KLT [3] and Parisi [7]. Starting from an initially flat interface above a d -dimensional substrate at time $t = 0$, a rough interface is described by its height $h(\vec{x}, t)$ at position $\vec{x}$ and time t , which obeys the following stochastic equation:

$$\frac{\partial}{\partial t} h(\vec{x}, t) = \nabla^2 h(\vec{x}, t) + F + \eta(\vec{x}, h), \quad (1)$$

where the diffusive term models the surface tension effects, F is the pushing force and the noise η is Gaussian,

with zero mean value and delta correlated,

$$\langle \eta(\vec{x}, y) \eta(\vec{x}', y') \rangle = \frac{\theta}{2} \delta^d(\vec{x} - \vec{x}') \delta(y - y'),$$

with an intensity θ . This equation also models the growth of domain walls in a random-field Ising ferromagnet [12].

Analytical results for the exponents associated with the model described by Eq. (1) are only known in the limit of strong pushing, where the interface moves very fast (for $d = 1$ the roughening exponent $\alpha = 1/2$ [3] and the time exponent $\beta = 1/4$ [7]). In this limit, the model reduces to the case of a noise with a delta correlation in time and the exponents correspond to the EW model, for which the scaling relation between exponents is given by $\alpha/\beta = 2$. On the other hand for small pushing ($F < F_c$) the interface becomes pinned.

KLT [3] and Parisi [7] have performed numerical simulations of the model given by Eq. (1) for different values of the pushing force. Far from F_c , Parisi [7] has obtained that, after a transient, there is a temporal regime with anomalous exponents followed by a crossover to an asymptotic regime that corresponds to the EW model. As one approaches the pinning transition, the crossover appears at later times, but the long-time behavior is described in any case by the EW model. Only at the pinning transition one gets a different exponent, $\beta = 3/4$ [7]. In the intermediate temporal regime, the numerical results of KLT [3] give $\alpha = 0.73$. These results should be compared with the experimental results of Rubio *et al.* [2] ($\alpha = 0.73 \pm 0.03$) and Horvath *et al.* [5] ($\beta = 0.65$ and $\alpha \sim 0.81$).

The purpose of this Letter is to obtain analytical expressions for the exponents characterizing the interface growth and the associated scaling relations in regimes where scaling properties are obeyed. To do so, we start with Eq. (1) and we derive an effective model valid for small intensities of the noise, θ . As we will see below, the resulting effective model is given by the following stochastic linear differential equation

$$\frac{\partial y}{\partial t} = \nabla^2 y + \xi(\vec{x}, t) \quad (2)$$

where $y(\vec{x}, t) = h(\vec{x}, t) - \bar{h}(t)$ measures the fluctuations of the height interface $h(\vec{x}, t)$ around its averaged value $\bar{h}(t)$. $\xi(\vec{x}, t)$ is the effective noise with zero mean value and correlation given by:

$$\langle \xi(\vec{x}, t) \xi(\vec{x}', t') \rangle = \frac{\theta}{2} \delta^d(\vec{x} - \vec{x}') W(t - t'). \quad (3)$$

The noise is delta-correlated in the substrate position and the dependence on h is substituted by a temporal correlation $W(t - t')$. The explicit form of $W(t - t')$ will be discussed in the following paragraph.

From Eqs. (2,3) it is possible to derive expressions for the width of the interface, $\sigma = \langle y^2 \rangle^{\frac{1}{2}}$ and the correlation length l_c in terms of $W(t - t')$. Therefore, the scaling properties of $W(t - t')$ will determine the roughening exponents and the associated scaling relations.

Derivation of the effective model.- In order to get an effective model, we use functional methods (a generalization [16] of the Novikov theorem [17]) to determine the mean value of η :

$$\begin{aligned} \langle \eta(\vec{x}, h(\vec{x}, t)) \rangle &= \frac{\theta}{2} \int_0^t G(0, t - t') dt' \times \\ &\times \int dy \int dy' \delta'(y - y') P_{\vec{x}, t; \vec{x}', t'}(y, y') + O(\theta^2) \end{aligned} \quad (4)$$

and also its correlation

$$\begin{aligned} \langle \eta(\vec{x}, h(\vec{x}, t)) \eta(\vec{x}', h(\vec{x}', t')) \rangle &= \frac{\theta}{2} \delta^d(\vec{x} - \vec{x}') \times \\ &\times \int dy P_{\vec{x}, t; \vec{x}', t'}(y, y) + O(\theta^2) \end{aligned} \quad (5)$$

where $G(\vec{x}, t) \sim t^{-d/2} \exp(-\frac{x^2}{t})$ is the diffusive propagator and

$$P_{\vec{x}, t; \vec{x}', t'}(y, y') = \langle \delta(y - h(\vec{x}, t)) \delta(y' - h(\vec{x}', t')) \rangle \quad (6)$$

is the joint probability for the interface to have heights y and y' at $\vec{x}, t$ and $\vec{x}', t'$ respectively. We note that Eqs.(4) and (5) are valid for small values of the intensity of the noise because we have neglected terms of order θ^2 involving the joint probability of four points. By assuming homogeneity both in space and time and defining $W(u, t - t')$ as the probability of having an increment u in a time $t - t'$, $W(u, t - t') = P_{\vec{x}, t; \vec{x}', t'}(u + h, h) / P_{\vec{x}, t'}(h)$, we obtain the effective model given by Eqs.(2,3) where $\xi(\vec{x}, t) = \eta(\vec{x}, h(\vec{x}, t)) - \langle \eta(\vec{x}, h(\vec{x}, t)) \rangle$ is the effective noise with zero mean value and correlation given by Eq.(5) up to the lowest order in θ , and $W(t - t') = W(0, t - t')$ is the probability for the interface to remain at the same point after a time interval $(t - t')$. Now, since the model is linear, the kinetics of the interface is completely characterized by this probability.

Exponents and scaling relations.- First, we will discuss the different relevant length scales associated with the interfacial growth. The interplay between these scales will determine the different scaling regimes. One scale is given by the horizontal correlation length, l_c , which for a very general form of $W(t - t')$, has diffusive behavior

with time, $l_c \sim t^{1/2}$. This result is remarkable because it indicates a very universal behavior for l_c in the complete temporal regime independent from the properties of the disorder. It only depends on the diffusive character of Eq.(2) and on the delta spatial correlation of the noise. Another relevant scale is associated to the characteristic length, L_c , of the clusters of points of the interface that become pinned by the effect of the quenched disorder. This cluster length depends on the pushing force. At the pinning transition, with a pushing force F_c , the entire interface becomes pinned and the cluster length is $L_c \sim L$. By increasing F , L_c is reduced and it becomes zero in the limit of strong pushing, where the interface moves very fast and the model reduces to EW model. In this limit, $W(t - t')$ becomes a delta function. In accordance with these results, we assume a scaling behavior of the form $L_c \sim (F - F_c)^{-\eta}$ where η is the associated exponent. Starting from a flat interface at $t = 0$ with a fixed value of the pushing force F , the horizontal correlation length, l_c , grows with time whereas the cluster length L_c , remains at a fixed value. At some time, that we define as the crossover time t_c , we have $l_c(t_c) = L_c$, and for times $t > t_c$ the evolution is described by the EW model. From its definition, t_c has the following scaling relation, $t_c \sim (F - F_c)^{-\nu}$, where $\nu = 2\eta$.

Regarding the short time behavior for which $l_c < L_c$, there is a regime with scaling properties in which the exponents and the scaling relations could be defined and they would be different from the ones associated with the EW model. To determine these quantities, we only need to realize that $W(t - t')$ depends on the relevant scale in this regime, L_c . However, due to the scaling properties, this is equivalent to assuming a dependence on L . In this way, we make a very general assumption about the form that $W(t - t')$ scales with L and t , $W(t - t') \sim L^{-\Omega} \sigma^{-\gamma}$ where Ω and γ are two independent exponents. Now, substituting this expression of $W(t - t')$ in the effective model, we obtain a scaling dependence of σ in L even before the saturation time is approached. Hence in this case the interface width scales with time and size before it saturates ($t < t_s$) as $\sigma \sim L^{\alpha'} t^{\beta}$ instead of the usual scaling given by $\sigma \sim t^{\beta}$. The expressions for the exponent β , the new exponent α' and the exponent z associated with l_c as $l_c \sim t^z$ are now given in terms of the parameters Ω and γ by:

$$\alpha' = -\frac{\Omega}{2 + \gamma} \quad \beta = \frac{4 - d}{2(2 + \gamma)} \quad z = 2 \quad (7)$$

From Eq.(7), we can discuss the expression of the α exponents that one gets for small system sizes for which the saturation happens before the effects of the pinning clusters have disappeared. The roughening exponent α is calculated by taking $t = t_s \sim L^2$ in the expression for $\sigma(t)$. The result is:

$$\alpha = \frac{4 - d}{2 + \gamma} - \frac{\Omega}{2 + \gamma} \quad (8)$$

A model for temporal correlation $W(t - t')$.- In the previous paragraph, we characterized the two different scaling regimes. Now, we would like to propose a simple model for $W(u, t - t')$ which match these two regimes. Based on physical arguments, one could assume that $W(u, t - t')$ is a regular function peaked on the mean height $u = \bar{h}(t - t')$ with a width σ . As a simple assumption, we consider a lorentzian function. In the short time behavior it has to have a scaling dependence in L and σ . This could simply be done by means of the following model

$$W(u, t) = \frac{\sigma}{\sigma^2 + (\bar{h} - u)^2} \left(L^\Omega \frac{\sigma^{1+\gamma}}{\bar{h}^2} + 1 \right)^{-1} \quad (9)$$

where γ and Ω are the parameters that characterized the model. The two different scaling regimes are defined by $L^\Omega \gg \bar{h}^2 \sigma^{-1-\gamma}$ and $L^\Omega \ll \bar{h}^2 \sigma^{-1-\gamma}$. From Eq. (9) we obtain that for long time, $t \gg t_c$ the scaling behavior of $W(0, t)$ is $W(0, \lambda t) \sim \lambda^{-1} \delta(t)$, as $\lambda \rightarrow \infty$, and $W(t - t')$ reduces to a delta function, as expected. By substituting $W(t - t')$ in the effective model one could get the exponents in the different scaling regimes. The advantage in assuming an expression for $W(t - t')$ over the complete temporal regime is that one can determine the crossover time, t_c , that can be defined by $L^\Omega = \bar{h}^2(t_c) \sigma^{-\gamma-1}(t_c)$. By substituting $\bar{h} \sim t$ and the scaling behavior of σ one obtains

$$t_c \sim L^{\Omega(2-\beta(1+\gamma))^{-1}(2+\gamma)^{-1}} \quad (10)$$

Numerical results and comparison.- At this point, all our exponents and scaling relations associated with the first scaling regime depend on the parameters Ω and γ . To corroborate the presence of both scaling regimes and the values of these parameters, we have performed numerical simulations of Eq.(1) for $d = 1$ with the same parameters as in Ref. [3]. In our numerical results, the pinning transition occurs at $F_c \sim 0.13$. We observe a first scaling regime with $\beta \sim 3/4$ in accordance with Parisi results. This result implies that our parameter γ is zero in this regime. After a crossover $t > t_c$, $\beta \sim 1/4$ like the EW model. In Fig. 1 we present our numerical results for l_c versus time in logarithmic scales. We observe the same diffusive behavior for the complete temporal regime in accordance with our model.

Regarding the roughening exponent α , our results are in agreement with those obtained by different methods, $\alpha = 0.7 - 0.9$ for $t_s < t_c$ [2-5,11] and $\alpha = 0.49$ for $t_s > t_c$ [5]. Furthermore, in the first regime, the results $\beta = 3/4$ and $\alpha = 0.8$ [18] are compatible in our analytical model by taking for our free parameters the values $\gamma = 0$ and $\Omega = 1.4$. Finally, in Fig. 2 we present our numerical results for the crossover time t_c versus $F - F_c$. We obtain an exponent $\nu \sim 1.43$, which in terms of L is equivalent to $t_c \sim L^{2-\nu} = L^{0.57}$. This numerical result is in good

agreement with our analytical results obtained by substituting in Eq.(10) the values of Ω , β and γ , i.e. $t_c \sim L^{0.56}$.

In conclusion, our analytical results predict a growth for the width in agreement with different simulations and experiments with the appropriate choice of two free parameters. A scaling relation determines the crossover time exponent. In our model, the correlation length shows a diffusive behavior and we believe it has to be analyzed in other models, like directed percolation (DP) and the random field Ising model, in order to clarify the diversity of growth behaviors and hence universality classes. For instance, in DP [9] anisotropy generates KPZ exponents after the crossover instead of those of EW.

The authors wish to acknowledge H. Guo for very fruitful discussions. This work has been supported by DGICYT of the Spanish Government, projects #PS90-0098 (J.M.L. and M.A.R.), #PB90-0030 (A.H.-M.), and #PB92-0863 (A.D.-G.), and NATO Collaborative Research Grant CRG.931018.

-
- [1] J. Koplik and H. Levine, Phys. Rev. B **32**, 280 (1985).
 - [2] M.A. Rubio, C.A. Edwards, A. Dougherty and J.P. Golub, Phys. Rev. Lett. **63**, 1685 (1989).
 - [3] D.A. Kessler, H. Levine and Y. Tu, Phys. Rev. A **43**, 4551 (1991).
 - [4] N. Martys, M. Cieplak and M.O. Robbins, Phys. Rev. Lett. **66**, 1058 (1991).
 - [5] V.K. Horvath, F. Family and T. Vicsek, J. Phys. A: Math. Gen. **24**, L25 (1991).
 - [6] T. Nattermann, S. Stepanow, L.-H. Tang and H. Leschhorn, J. Phys. II France **2**, 1483 (1992).
 - [7] G. Parisi, Europhys. Lett. **17**, 673 (1992).
 - [8] K. Sneppen, Phys. Rev. Lett. **69**, 3539 (1992).
 - [9] L.-H. Tang and H. Leschhorn, Phys. Rev. A **45**, R8305 (1992).
 - [10] Z. Csahok, K. Honda and T. Vicsek, J. Phys. A: Math. Gen. **26**, L171 (1993).
 - [11] C.S. Nolle, B. Koiller, N. Martys and M.O. Robbins, Phys. Rev. Lett. **71**, 2074 (1993).
 - [12] O. Narayan and D.S. Fisher, Phys. Rev. B **48**, 7030 (1993).
 - [13] S.F. Edwards and D.R. Wilkinson, Proc. R. Soc. Lond. **A381**, 17 (1982).
 - [14] M. Kardar, G. Parisi and Y.-C. Zhang, Phys. Rev. Lett. **56**, 889 (1986).
 - [15] F. Family T. Vicsek eds., *Dynamics of fractal surfaces*, (World Scientific, 1991, Singapore).
 - [16] P. Hanggi, in *Stochastic processes applied to physics*, L. Pesquera and M. A. Rodriguez eds., (World Scientific, 1985, Singapore) p.69.
 - [17] E.A. Novikov, Sov. Phys. JETP **20**, 1290 (1965).
 - [18] The range of values for α obtained in different works seems to be due to a different interpretation of overhangs as is discussed in [11]. Our model (1) does not contain

overhangs because the partial differential equation has a single-valued solution.

FIG. 1. Horizontal correlation length versus time for two values of F , close $F = 0.15$ and far $F = 0.4$ from the pinning transition $F_c = 0.13$. The slope of the adjusted curve is $1/2$.

FIG. 2. Crossover time versus $F - F_c$ in log-log scale. The adjusted curve gives an exponent $\nu = 1.43$.